

/*05LABOKLIN NV . Verlengde Klinkertstraat 6 . NL-6433PL Hoensbroek/*02

Centre Vet 484 Chaussée de Bruxelles, 484 7850 Enghien Belgien

/*05Report/*14 No.: 1806-N-05872

Date of arrival: 05-06-2018 Date of report: 12-06-2018

Patient identification: dog male * 06.03.17

Pug

Owner / Animal-ID: Montens, Christian

Type of sample: EDTA

Date sample was taken: 01-06-2018

Name: Arthur's Excalibur Rising By Megipugi

Stud book no.: AV 01127501 Chip no.: 981000010206908

Tattoo no.:

Necrotizing Meningoencephalitis (PDE) - PCR

Result: Genotype N/N

Interpretation: The examined animal is homozygous for the wildtype-allele. It does not carry the high-risk factor for NME in the DLA-DPB1-gene.

Trait of inheritance: autosomal-recessive

Scientific studies found correlation between the mutation and symptoms of the disease in the following breeds: Pug

Degenerative Myelopathy - PCR

Result: Genotype N/N (exon 2)

Interpretation: The examined animal is homozygous for the wildtype-allele. It does not carry the high-risk factor for DM in exon 2 of the SOD1-gene.

sample ID: 1806-N-05872



Trait of inheritance: autosomal-recessive

Please note: In the Bernese Mountain Dog breed the mutation in exon 1 of the SOD1-gene also occurs in correlation with DM.

The current result is only valid for the sample submitted to our laboratory. The sender is responsible for the correct information regarding the sample material. The laboratory can not be made liable. Furthermore, any obligation for compensation is limited to the value of the tests performed.

There is a possibility that other mutations may have caused the disease/phenotype. The analysis was performed according to the latest knowledge and technology.

The laboratory is accredited for the performed tests according to DIN EN ISO/IEC 17025:2005. (except partner lab tests).

*** END of report ***

Drs J. Vis

* * * It's breeding season again * * *

Bacteriological testing of cervical swabs offers important information on the clinical health of mares. Bacterial culture and identification of isolates and resistance testing by microdilution help in treating acute and subclinical infections.